

ROOT NODULE BACTERIA
STORE IN A COOL PLACE

Nodulaid[®]

A potent inoculant for nitrogen fixation and increased yields

GROUP J

Pigeon Pea, Lablab

For inoculating up to 100kg of Pigeon Pea and Lablab and up to 50kg of Axillaris seed.
Refer to direction of use for Groups E, F, G, N & H

BASF Australia Ltd ABN 62 008 437 867
Level 12, 28 Freshwater Place Southbank VICTORIA 3006, AUSTRALIA
Website: www.agro.basf.com.au

BASF New Zealand Ltd 110297
3 Airpark Drive, Airport Oaks, MANKAU 2022, NEW ZEALAND
Website: www.agro.basf.com.nz

Customer Service Hotline: 1800 558 399

® Registered trademark of BASF Agricultural Specialities Pty Ltd

DIRECTIONS FOR USE with groups AL, B, C**SLURRY INOCULATION**

1. Prepare a solution of BASF Seedstik coating adhesive.
2. Prepare a slurry by mixing the BASF Seedstik solution with the entire contents of this pack and 5L of cool, clean water.
3. Stir to ensure inoculant is thoroughly dispersed.
4. Pour slurry over the correct weight of seed and mix until all the seeds are well coated. Mixing can be done in clean concrete mixer or similar vessel, or added to the seed during transfer stages e.g. auguring seed.
5. Sow inoculated seed as soon as possible or within 24 hours.

DIRECTIONS FOR USE with Group's E, F, G, N & H**SLURRY INOCULATION**

1. Prepare a slurry by mixing the entire contents of this pack into 5 litres of cool, clean water.
2. Stir to ensure inoculant is thoroughly dispersed. Wait for a minimum of 15 minutes, stirring occasionally to allow inoculant adhesive to dissolve.
3. Pour slurry over the correct weight of seed and mix until all seeds are wet. Mixing can be done in a clean concrete mixer or similar vessel, or added to seed during transfer stages, e.g. auguring seed.
4. Use the slurry mix within 24 hours. Sow inoculated seed as soon as possible or store for up to 3 days in a COOL place, away from sunlight.

FURROW APPLICATION INSTRUCTIONS

1. Prepare in furrow solution directly in a clean liquid injection cart.
2. Fill cart with cool, clean water and either add the Nodulaid directly to the water and mix well or place the Nodulaid into a stocking and suspend inside the tank.
3. Apply at 50-100L/Ha depending on row spacing.
4. Use solution within 24hrs.
5. Ensure coarse nozzles are used to avoid blockages.

PRECAUTIONS

Store this pack of inoculant in a COOL place. Do not store opened packs of inoculant. Do not use out-of-date inoculants. Avoid contact with the eyes - may cause irritation. Avoid inhaling dust - may cause irritation.

PESTICIDES

Most pesticides are toxic to inoculants. Do not use any vessel which has contained poisonous sprays or dusts. Check with your State Department or Agriculture or BASF for compatibility.

FERTILIZERS

Slurry inoculated seed should never be mixed with very cold or alkaline fertilizer which will KILL the inoculant bacteria. Excessive use of starter Nitrogen fertilizer, or high levels of soil nitrate can delay or reduce nodulation.

WARRANTY

The product must be used strictly in accordance with the label directions and instructions. BASF accepts no liability or responsibility for any consequential loss or otherwise arising from failure to follow these directions and instructions. The liability of BASF is limited to that imposed by Statutes and, in any case, the liability is strictly limited to replacement of the product or (at the option of BASF) the refund of the price paid.

Batch No:

Date of Manufacture:

BASF Australia Ltd
ABN 62 008 437 867
Level 12, 28 Freshwater Place
Southbank VICTORIA 3006
AUSTRALIA

FOR SPECIALIST ADVICE IN AN EMERGENCY ONLY PHONE 1800 803 440 TOLL FREE – ALL HOURS – AUSTRALIA WIDE

BASF New Zealand Ltd
110297
3 Airpark Drive,
Airport Oaks, MANKAU 2022,
NEW ZEALAND

EMERGENCY PHONELINK 0800944 955 TOLL FREE – ALL HOURS – NEW ZEALAND WIDE