

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

# GREEN GUARD<sup>®</sup> SC BIOLOGICAL INSECTICIDE

ACTIVE CONSTITUENT: *Metarhizium anisopliae* var. *acridum* spores (4 x 10<sup>10</sup>/gram)

For the Biological Control of Locusts and Grasshoppers

This pack contains 50 g of *M. anisopliae* var. *acridum* spores in a plastic bag, 250mL surfactant solution in an inner pouch and 300 mL emulsifiable oil in a bottle (to be mixed as per Mixing Instructions).

The bag, pouch and bottle must not be sold separately.  
Do not destroy this pack while inner contents still remain.

CONTENTS: 600g, 1.8kg

BASF Australia Ltd ABN 62 008 437 867  
Level 12, 28 Freshwater Place Southbank VICTORIA 3006, AUSTRALIA  
Website: [www.agro.basf.com.au](http://www.agro.basf.com.au)

APVMA Approval No: 55736/600g/0905, 55736/1.8kg/0905

Customer Service Hotline: 1800 558 399

® Registered trademark of CSIRO



## DIRECTIONS FOR USE

### RESTRAINTS

DO NOT proceed to use this pack until all instructions are read carefully.

Mixing incorrectly will reduce the effectiveness of the product.

DO NOT apply in gusty conditions greater than 8 metres per second or if rain is imminent within 6 hours.

DO NOT apply in gusty conditions greater than 8 meters per second or if rain is imminent within 6 hours.

SITUATION	PEST	RATE/ha	CRITICAL COMMENTS
Agricultural areas, Pastures, Crops, Forage crops and Non-crop areas	Nymphs of the Australian plague locust ( <i>Chortoicetes terminifera</i> ), Wingless grasshopper ( <i>Phaulacridium vittatum</i> ) and pest grasshoppers	75 g spores (the final mixed product diluted into 150 – 450 L of water treats 2 hectares)	The supplied product contains three different components and requires mixing prior to use*. For best results, apply when locusts and grasshoppers are at early nymph stage. Do NOT apply in gusty conditions greater than 8 metres per second or if rain is imminent in the next 6 hours. *Refer to “Mixing Instructions” below; incorrect mixing will reduce the effectiveness of the product.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

### GENERAL INSTRUCTIONS

The pail contains: 1) a bottle of 900 mL emulsifiable oil and, 2) a clear plastic bag containing 150 g of *Metarhizium anisopliae* var. *acidum* dry spores plus 750 mL of surfactant solution in a silver inner pouch. The inner pouch inside the clear plastic bag may not be readily visible if covered with spores. **Read “Mixing Instructions” before opening the bag or bottle.** GREEN GUARD® SC is specifically designed for small-scale application in agricultural areas, pastures, crops, forage crops and non-crop areas. At the warm to hot (25° – 35°C) temperatures common during spring and summer, locusts and grasshoppers treated with GREEN GUARD® SC take 8-12 days to die. Mortality may take longer when it is cool or cloudy. Because mortality can take 8-12 days or more, the best strategy with most locusts and grasshoppers is to begin treatment early in the nymphal stage. When locusts or grasshoppers are spread out, treat all of the infested area. However, locusts sometimes form discrete bands. In such cases, first treat the vegetation for 10-15 m in front of the band and then treat the band itself. The nymphs will not only be treated directly but will also pick up a further dose from the vegetation as they march through it.

### MIXING INSTRUCTIONS

1. Remove the clear plastic bag and the bottle from the pail. Do NOT open the clear plastic bag or the bottle until instructed to do so.
2. Lay the clear plastic bag on a firm surface then locate the silver inner pouch within the clear plastic bag. With a clenched fist, strike to break the seal of the inner pouch, the seal is designed to break upon impact and release the surfactant solution still within the clear plastic bag.
3. Once the seal of the inner pouch is broken, thoroughly mix the surfactant solution and dry spores in the unopened clear plastic bag for 2 - 3 minutes. Let it stand 10 minutes to allow the spores to be well coated with surfactant solution.
4. Tear open the clear plastic bag and pour the mixture of spores and surfactant solution into the pail. Use a small amount of water to rinse the spore residues and add rinsings to the pail. Dispose of the empty bag and pouch accordingly (refer Storage and Disposal section).
5. Add the entire contents of the bottle containing 900 mL of emulsifiable oil into the pail, mix gently with stirring stick supplied until oil is dispersed evenly throughout solution (approximately 1 minute). Allow to sit for 2-3 minutes to allow spores and oil to come together. Rinse the bottle and add rinsings to the pail. Disposal of empty bottle accordingly (refer Storage and Disposal section).
6. Add the mixed solution into a clean spray tank and dilute with drinking quality water to make 150-450L of spray. Use

more water (400-450 L) for tall and thick vegetation. Triple or preferably pressure rinse the pail before disposal. Add rinsings to spray tank. Dispose of the empty pail accordingly (refer Storage and Disposal section).

7. Mix thoroughly. To ensure product effectiveness, diluted product should be used immediately.

### APPLICATION

Ensure equipment being used is free of chemical residues. The formulation is designed to be applied through the diverse range of nozzle or boom spray equipment commonly used to apply water-based sprays. Apply by ground based equipment to achieve a droplet size (VMD) in the range 80-250 µm. As the active ingredient consists of small particles, very fine filters will sometimes clog up. If there is clogging, remove the filter and replace with a filter with a larger mesh. Apply when temperature is between 15° – 30°C.

### COMPATIBILITY

This product should not be used with other pesticides.

### PRECAUTIONS

**Re-entry periods:** DO NOT enter treated area for at least 24 hours after spraying, unless wearing a disposable mist mask, cotton overalls buttoned to neck and wrist and washable hat and rubber gloves. After each day's use, wash clothing and gloves.

### PROTECTION OF BEES

This product is a potential hazard to bees. DO NOT spray on plants in flower while bees are foraging.

### PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with GREEN GUARD® SC or with used containers. DO NOT apply under weather conditions, or from spray equipment, that may cause spray to drift onto nearby natural streams, rivers or waterways. DO NOT apply within 5 m upwind of wetlands, waterbodies or watercourses.

### STORAGE AND DISPOSAL

Keep out of reach of children. Store in the closed, original container below 5°C (do not freeze). Just prior to use, short term storage at room temperature is acceptable. DO NOT store in direct sunlight. Triple or preferably pressure rinse empty pails and rinse empty bags and bottles in water before disposal. Add rinsings to spray tank. DO NOT dispose undiluted product on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

### SAFETY DIRECTIONS

Avoid contact with eyes. May irritate the eyes. Repeated exposure may cause allergic disorders. Sensitive workers should use protective clothing. When preparing the product for use, wear cotton overalls buttoned to neck and wrist and a washable hat, elbow-length rubber gloves and face shield. When using the prepared spray by other than aerial application, wear a disposable mist mask. After use, wash hands thoroughly with soap and water.

### FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre (Phone Australia 131126).

### MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet.

### WARRANTY

The product must be used strictly in accordance with the label directions and instructions. BASF accepts no liability or responsibility for any consequential loss or otherwise arising from failure to follow these directions and instructions. The liability of BASF is limited to that imposed by Statutes and, in any case, the liability is strictly limited to replacement of the product or (at the option of BASF) the refund of the price paid.

® GREEN GUARD is a registered trade mark of the Commonwealth Scientific and Industrial Research Organisation (CSIRO).

APVMA Approval No. 55736/600g/0905, 55736/1.8kg/0905

Batch No:

Date of Manufacture:

BASF Australia Ltd  
ABN 62 008 437 867  
Level 12, 28 Freshwater Place  
Southbank VICTORIA 3006  
AUSTRALIA

FOR SPECIALIST ADVICE IN AN EMERGENCY ONLY PHONE 1800 803 440 TOLL FREE – ALL HOURS – AUSTRALIA WIDE