



CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

PREMIS® PRO C SEED TREATMENT

**ACTIVE CONSTITUENT: 25 g/L TRITICONAZOLE
4 g/L CYPERMETHRIN**

GROUP	3	FUNGICIDE
GROUP	3A	INSECTICIDE

For the control of covered smut, flag smut and loose smut in wheat, covered smut and loose smut in barley and oats, and for protection against insect pests of stored grain.

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USING THIS PRODUCT

CONTENTS: 10L-100L

BASF Australia Ltd ABN 62 008 437 867
Level 12, 28 Freshwater Place Southbank VICTORIA 3006

® Registered trademark of BASF

APVMA Approval No.: 58965/56990



STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

10L

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

100L

If tamper evident seals are broken prior to initial use then the integrity of the contents cannot be assured. Empty container by pumping through dry-break connection system. Do not attempt to breach the valve system or the filling point, or contaminate the container with water or other products. Ensure that the coupler, pump, meter and hoses are disconnected, triple rinsed and drained after each use. When empty, or contents no longer required, return the container to the point of purchase. This container remains the property of BASF.

SAFETY DIRECTIONS

Will irritate the eyes and skin. Avoid contact with eyes and skin. Facial skin contact may cause temporary facial numbness. When opening the container and preparing the product for use wear elbow-length PVC gloves. Wash hands after use. After each day's use wash gloves.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 131126.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet.

CONDITIONS OF SALE

All conditions and warranties rights and remedies implied by law or arising in contract or tort whether due to the negligence of BASF Australia Ltd or otherwise are hereby expressly excluded so far as the same may legally be done provided however that any rights of the Buyer pursuant to non-excludable conditions or warranties of the Trade Practices Act 1974 or any relevant legislation of any State are expressly preserved but the liability of BASF Australia Ltd or any intermediate Seller pursuant thereto shall be limited if so permitted by the said legislation to the replacement of the goods sold or the supply of equivalent goods and all liability for indirect or consequential loss or damage of whatsoever nature is expressly excluded. This product must be used or applied strictly in accordance with the instructions appearing hereon. This product is solely sold for use in Australia and must not be exported without the prior written consent of BASF Australia Ltd.

APVMA Approval No: 58965/56990

Batch No:

Date of Manufacture:

BASF Australia Ltd
ABN 62 008 437 867
Level 12, 28 Freshwater Place
Southbank VICTORIA 3006

FOR SPECIALIST ADVICE IN AN EMERGENCY ONLY PHONE 1800 803 440 TOLL FREE-ALL HOURS-AUSTRALIA WIDE

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

PREMIS® PRO C SEED TREATMENT

**ACTIVE CONSTITUENT: 25 g/L TRITICONAZOLE
4 g/L CYPERMETHRIN**

GROUP	3	FUNGICIDE
GROUP	3A	INSECTICIDE

For the control of covered smut, flag smut and loose smut in wheat, covered smut and loose smut in barley and oats, and for protection against insect pests of stored grain.

IMPORTANT: READ THIS LEAFLET BEFORE USING THIS PRODUCT

CONTENTS: 10-100 L

BASF Australia Ltd ABN 62 008 437 867
Level 12, 28 Freshwater Place Southbank VICTORIA 3006

® Registered trademark of BASF

APVMA Approval No.: 58965/56990

DIRECTIONS FOR USE

CROP	DISEASE/PEST	RATE PER 100 kg SEED	CRITICAL COMMENTS
Wheat	Covered smut (Bunt) (<i>Tilletia</i> spp.), Flag smut (<i>Urocystis agropyri</i>), Loose smut (<i>Ustilago tritici</i>)	100 mL	Ensure even and thorough coverage of the seed. Use of 400 mL of mixture (water + product) with each 100 kg seed is recommended.
Barley	Covered smut (<i>Ustilago segetum</i> var. <i>hordei</i>), Loose smut (<i>Ustilago tritici</i>)		
Oats	Covered smut (<i>Ustilago segetum</i> var. <i>hordei</i>), Loose smut (<i>Ustilago avenae</i>)		
Wheat, barley, oats	Protection against insect pests of stored grains: Lesser grain borer, granary weevil, rice weevil, rust-red flour beetle, saw toothed grain beetle, tropical warehouse moth		

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:

WHEAT, BARLEY and OATS: NOT REQUIRED WHEN USED AS DIRECTED.

GENERAL INSTRUCTIONS

Protection from smuts and bunts will only be achieved if all grains are treated. Therefore ensure that the application equipment is correctly set up to achieve this. In Queensland, the emergence of treated seed sown too deep or into warm soil may be affected. Sowing rates may need to be adjusted accordingly.

FUNGICIDE RESISTANCE WARNING

GROUP 3 FUNGICIDE

Premis Pro C Seed Treatment is a member of the DMI group of fungicides. For fungicide resistance management the product is a Group 3 fungicide. Some naturally occurring individual fungi resistant to the product and other Group 3 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by Premis Pro C and other Group 3 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, BASF Australia Ltd accepts no liability for any losses that may result from the failure of Premis Pro C to control resistant fungi.

INSECTICIDE RESISTANCE WARNING

GROUP 3A INSECTICIDE

For insecticide resistance management, Premis Pro C Seed Treatment is a Group 3A insecticide. Some naturally occurring insect biotypes resistant to Premis Pro C and other Group 3A insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Premis Pro C or other Group 3A insecticides are used repeatedly. The effectiveness of Premis Pro C on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, BASF Australia Ltd accepts no liability for any losses that may result from the failure of Premis Pro C to control resistant insects. Premis Pro C may be subject to specific resistance management strategies. For further information contact your local supplier, BASF representative or local agricultural department agronomist.

Mixing

Shake the container well before using. Add the required volume of Premis Pro C to the water whilst agitating.

Application

Ensure the application thoroughly covers the seed. For information about suitable application equipment and calibration, contact your supplier of Premis Pro C or a BASF representative. Wash equipment thoroughly with water after application.

Compatibility

Do not mix with any other product.

PRECAUTIONS

When treated seed is stored, it should be kept apart from other grain or seeds, and the containers or bags should be clearly marked to show that the contents have been treated with Premis Pro C Seed Treatment.

DO NOT use treated seed for human consumption.

DO NOT allow treated seed to contaminate food for human consumption.

Bags which have held treated seed should not be used for any other purpose. Any seed not used for sowing should be destroyed.

PROTECTION OF LIVESTOCK

DO NOT feed treated seed to animals, including poultry.

Store treated seed away from feed.

DO NOT allow treated seed to contaminate grain intended for animal consumption.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS, AND ENVIRONMENT

DO NOT feed treated seed or otherwise expose to wildlife or domestic birds. DO NOT contaminate streams, rivers or waterways with the chemical, used containers, treated seed or bags which have held treated seed.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

10L

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

100L

If tamper evident seals are broken prior to initial use then the integrity of the contents cannot be assured. Empty container by pumping through dry-break connection system. Do not attempt to breach the valve system or the filling point, or contaminate the container with water or other products. Ensure that the coupler, pump, meter and hoses are disconnected, triple rinsed and drained after each use. When empty, or contents no longer required, return the container to the point of purchase. This container remains the property of BASF.

SAFETY DIRECTIONS

Will irritate the eyes and skin. Avoid contact with eyes and skin. Facial skin contact may cause temporary facial numbness. When opening the container and preparing the product for use wear elbow-length PVC gloves. Wash hands after use. After each day's use wash gloves.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 131126.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet.

CONDITIONS OF SALE

All conditions and warranties rights and remedies implied by law or arising in contract or tort whether due to the

negligence of BASF Australia Ltd or otherwise are hereby expressly excluded so far as the same may legally be done provided however that any rights of the Buyer pursuant to non-excludable conditions or warranties of the Trade Practices Act 1974 or any relevant legislation of any State are expressly preserved but the liability of BASF Australia Ltd or any intermediate Seller pursuant thereto shall be limited if so permitted by the said legislation to the replacement of the goods sold or the supply of equivalent goods and all liability for indirect or consequential loss or damage of whatsoever nature is expressly excluded. This product must be used or applied strictly in accordance with the instructions appearing hereon. This product is solely sold for use in Australia and must not be exported without the prior written consent of BASF Australia Ltd.

APVMA Approval No: 58965/56990

Batch No:

Date of Manufacture:

® = Registered trademark of BASF



The Chemical Company

BASF Australia Ltd
ABN 62 008 437 867
Level 12, 28 Freshwater Place
Southbank VICTORIA 3006

FOR SPECIALIST ADVICE IN AN EMERGENCY ONLY PHONE 1800 803 440 TOLL FREE-ALL HOURS-AUSTRALIA WIDE.