



CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

SYSTIVA® SEED TREATMENT FUNGICIDE

ACTIVE CONSTITUENT: 333 g/L FLUXAPYROXAD

GROUP	7	FUNGICIDE
-------	----------	-----------

For the control of a range of seed and foliar fungal diseases of barley and wheat and the suppression of rhizoctonia root rot in barley and wheat.

CONTENTS: 1- 20 L

BASF Australia Ltd ABN 62 008 437 867
Level 12, 28 Freshwater Place Southbank VICTORIA 3006

® Registered trademark of BASF

APVMA Approval No.: 70065/61412

DIRECTIONS FOR USE

Crop	Disease	Application Rate	Critical Comments
Barley	Loose smut (<i>Ustilago segetum</i> var <i>nuda</i>) Suppression of Rhizoctonia root rot (<i>Rhizoctonia</i> <i>solanii</i>)	150mL/100kg seed	Apply diluted with sufficient water to ensure even coverage of seed. Apply with an applicator designed for liquid seed treatments. Calibrate application equipment for the flow rate of the grain. SYSTIVA will provide suppression of Rhizoctonia and reduce severity of disease within the crop.
	Powdery Mildew (<i>Blumeria</i> <i>graminis</i> f.sp. <i>hordei</i>) Net Form of Net Blotch (<i>Pyrenophora</i> <i>teres</i> f. <i>teres</i>) Scald (<i>Rhynchosporium</i> <i>secalis</i>) Spot form of Net Blotch (<i>Pyrenophora</i> <i>teres</i> f. <i>maculata</i>) Leaf rust (<i>Puccinia hordei</i>)		SYSTIVA is systemic and will provide ongoing residual activity. Typically, greater control can be seen on newly emerged, younger leaves. Foliar diseases will be controlled in the crop canopy, including upper leaves, up to the time when the crop reaches the following growth stages,; Powdery Mildew and Leaf Rust will be controlled up to flag leaf emergence – BBCH 37. Net Form of Net Blotch, Spot Form of Net Blotch and Scald will be controlled up to at least the middle of heading – BBCH 55 (middle of inflorescence emerging),
Wheat	Bunt (<i>Tilletia</i> spp.) Suppression of Rhizoctonia root rot (<i>Rhizoctonia</i> <i>solanii</i>)	150mL/100kg seed	Applied diluted with sufficient water to ensure even coverage of seed. Apply with an applicator designed for liquid seed treatments. Calibrate application equipment for the flow rate of the grain. SYSTIVA will provide suppression of Rhizoctonia and reduce severity of disease within the crop.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:

GRAZING: DO NOT GRAZE PLANTS GROWN FROM TREATED SEED OR CUT FOR STOCK FOOD WITHIN 4 WEEKS OF SOWING

HARVEST: NOT REQUIRED WHEN USED AS DIRECTED

LIVESTOCK DESTINED FOR EXPORT MARKETS

The grazing withholding period only applies to stock slaughtered for the domestic market. Some export markets apply different standards. To meet these standards, ensure that in addition to complying with the grazing withholding period, the Export Slaughter Interval is observed before stock are sold or slaughtered.

EXPORT SLAUGHTER INTERVAL (ESI)

After observing the withholding period for grazing or cutting for stockfood, livestock that have been grazed on or fed treated crops should be placed on clean feed for 2 days prior to slaughter.

EXPORT OF TREATED CEREALS

Growers should note that Maximum Residue Limits (MRLs) or import tolerances do not exist in all markets for cereals treated with SYSTIVA. Additionally, some export markets have established MRLs different to those in Australia. If you are growing cereals for export, please check with BASF or your grain exporter for the latest information on MRLs and import tolerances BEFORE using SYSTIVA.

GENERAL INSTRUCTIONS

SYSTIVA is a broad spectrum systemic fungicide. It inhibits spores germination, mycelial growth and sporulation of the fungus. Application of SYSTIVA to seed will result in control of seed borne loose smut in barley and bunt in wheat, suppression of Rhizoctonia and extended control of foliar diseases as outlined in the Directions for Use table. SYSTIVA treated seed will act as a reservoir and continue to provide systemic movement to the foliage over an extended period of time.

APPLICATION

Apply to seed with equipment which results in even coverage of all seed. SYSTIVA should be mixed with a sufficient volume of water up to a maximum mix volume of 600mL/100kg seed, to ensure a slurry which results in thorough and even coverage of seed.

Storage of Treated Seed

Seed treated with SYSTIVA may be stored in a cool, dry well ventilated place until the following season, with no adverse effects on germination or emergence, however as conditions of storage may not be known, BASF accept no liability for stored treated seed.

When treated seed is stored, it should be kept apart from other grain and the bags or containers should be clearly marked to indicate the contents have been treated with this product. DO NOT use treated seed for human consumption. Bags which have held treated seed are not to be used for any other purpose.

FUNGICIDE RESISTANCE WARNING

GROUP	7	FUNGICIDE
--------------	----------	------------------

SYSTIVA is a member of the succinate dehydrogenase inhibitor (SDHI) group of fungicides. For fungicide resistance management the product is a Group 7 fungicide.

Some naturally occurring individual fungi resistant to SYSTIVA and other Group 7 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if SYSTIVA is used repeatedly. These resistant fungi may not be controlled by SYSTIVA or other Group 7 fungicides, thus resulting in a reduction in efficacy and possible yield loss.

Since the occurrence of resistant fungi is difficult to detect prior to use, BASF Australia Ltd accepts no liability for any losses that may result from the failure of SYSTIVA to control resistant fungi.

RE-HANDLING TREATED SEED

Do not allow re-handling of treated seed until dry unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

PRECAUTIONS

Do not use treated seed for human or animal consumption. Do not allow treated seed to contaminate seed intended for human or animal consumption.

PROTECTION OF LIVESTOCK

DO NOT feed treated seed to animals, including poultry. DO NOT allow treated seed to contaminate grain intended for animal consumption.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT feed treated seed or otherwise expose to wild or domestic birds. Any spillages of treated seed, however minor, must be cleaned up immediately, preferably by recovery and re-use. If disposal is required ensure treated seeds are thoroughly buried and not accessible to birds and other wildlife.

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area out of direct sunlight. Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. DO NOT burn empty containers or product.

SAFETY DIRECTIONS

May irritate the skin. Avoid contact with skin. When opening the container and mixing and loading and using the product, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow-length chemical resistant gloves. Wash hands after use. After each day's use wash gloves and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet available from www.basf.com.au.

CONDITIONS OF SALE

All conditions and warranties rights and remedies implied by law or arising in contract or tort whether due to the negligence of BASF Australia Ltd or otherwise are hereby expressly excluded so far as the same may legally be done provided however that any rights of the Buyer pursuant to non-excludable conditions or warranties of the Trade Practices Act 1974 or any relevant legislation of any State are expressly preserved but the liability of BASF Australia Ltd or any intermediate Seller pursuant thereto shall be limited if so permitted by the said legislation to the replacement of the goods sold or the supply of equivalent goods and all liability for indirect or consequential loss or damage of whatsoever nature is expressly excluded. This product must be used or applied strictly in accordance with the instructions appearing hereon. This product is solely sold for use in Australia and must not be exported without the prior written consent of BASF Australia Ltd.

APVMA Approval No: 70065/61412

Batch No:

Date of Manufacture:

® = Registered trademark of BASF



The Chemical Company

BASF Australia Ltd

ABN 62 008 437 867

Level 12, 28 Freshwater Place

Southbank VICTORIA 3006

FOR SPECIALIST ADVICE IN AN EMERGENCY ONLY PHONE 1800 803 440 TOLL FREE-ALL HOURS-AUSTRALIA WIDE.